

[preprint]aastex

12 November 1999

To appear in 2000 Apr 10 issue of ApJ.

χ^2 et al. $\text{km s}^{-1} \sim$ *HEAO-1 ASCA Ginga ROSAT EXOSAT Einstein BeppoSAX* X1832–330
NGC 6652 Paper I

LMXB in NGC 6652 Mukai & Smale

document

The Low-Mass X-ray Binary X1832–330 in the Globular Cluster NGC 6652: A Serendipitous ASCA
Observation

Koji Mukai¹ and Alan P. Smale¹ Code 662, NASA/Goddard Space Flight Center, Greenbelt, MD 20771,
USA.

¹Also Universities Space Research Association

abstract

The Low Mass X-ray Binary (LMXB) in is one of 12 bright, or transient, X-ray sources to have been discovered in Globular Clusters. We report on a serendipitous *ASCA* observation of this Globular Cluster LMXB, during which a Type I burst was detected and the persistent, non-burst emission of the source was at its brightest level recorded to date. No orbital modulation was detected, which argues against a high inclination for the system. The spectrum of the persistent emission can be fit with a power law plus a partial covering absorber, although other models are not ruled out. Our time-resolved spectral analysis through the burst shows, for the first time, clear evidence for spectral cooling from $kT=2.4\pm0.6$ keV to $kT=1.0\pm0.1$ keV during the decay. The measured peak flux during the burst is $\sim 10\%$ of the Eddington luminosity for a $1.4M_{\odot}$ neutron star. These are characteristic of a Type I burst, in the context of the relatively low quiescent luminosity of .